

POST-OPERATIVE INSTRUCTIONS FOR THE CARE OF YOUR CHILD

Your child has received one or more of the following circled items. Please follow the instructions below for his/her proper care.

If you have any questions, please call Dr Grant at 301-731-8510 or email drgrant@mychildrenschoicedental.com.

LOCAL ANESTHETIC	<ul style="list-style-type: none"> ◆Your child has been given an injection of local anesthetic to numb his/her teeth in order for us to work on them. Your child's mouth (tooth, lip & tongue) will be sleeping for approximately 2 to 3 hours. It is imperative that you watch your child and have him/her bite on the cotton roll provided to prevent damage to his/her oral cavity. If your child is not biting on the cotton roll he/she may bite the tongue, cheek or lip causing serious damage or profound pain later on. ◆If your child falls asleep you may remove the cotton roll to prevent him/her from swallowing or choking on it. ◆Some children become very upset (even crying hysterically) and complain of excruciating pain when they realize their mouth feels "different." Please do not be alarmed! Many children are unfamiliar with this numb sensation and associate this with pain. Reassure your child that the "funny feeling" will go away in approximately 2 hours.
CLEANING	<ul style="list-style-type: none"> ◆Your child has received a dental a cleaning which included the application of fluoride. The fluoride is applied to the teeth with a tooth brush after the cleaning in an effort to prevent cavities from developing. Because of this it is important that you not allow your child to eat or drink anything for 20 minutes to enable the fluoride to "soak in." The application of fluoride is recommended every 6 months.
SHAVE	<ul style="list-style-type: none"> ◆Shaving of enamel is indicated in the following situations: (1) to prevent a very small cavity from enlarging (2) to correct an ectopic eruption (3) to correct mild crowding (interceptive orthodontics)
SEALANT	<ul style="list-style-type: none"> ◆A sealant has been placed on one or more of your child's teeth in an effort to prevent him/her from developing a cavity on the top surface of the tooth. Your child must avoid eating ice or chewing on hard sticky candies which can dislodge or chip the sealant, rendering it ineffective. Please also be advised that sealants do not protect the surfaces in-between teeth so your child must still floss to prevent this type of cavity from developing between the teeth.
FILLING	<ul style="list-style-type: none"> ◆Most children usually adapt and adjust very well to new dental work. However, if some fillings are a bit deep due to the extent of the decay, they may complain of some discomfort. Do not be alarmed! Sometimes it takes a few weeks for the tooth to settle down and feel comfortable.
CROWN (SILVER OR WHITE)	<ul style="list-style-type: none"> ◆If your child has received a crown/cap the gum tissue surrounding the tooth may appear to be bleeding. This is normal! When preparing your child's tooth for a crown, it is necessary to fit the crown closely to the gum tissue, resulting in some hemorrhaging and bruising of the tissue. Following placement of the crowns some parents will notice a purple or gray color around the gums; this is also normal and will subside. ◆Please do not allow your child to eat sticky or caramel type candy as this may dislodge your child's crown. This applies for as long as the crown is in your child's mouth. ◆If your child's crown does come off please save the crown and call the office so it may be recemented.
NERVE TREATMENT	<ul style="list-style-type: none"> ◆Please note that children seldom complain of pain after this procedure is performed. When your child receives a nerve treatment it is usually not necessary to prescribe any pain medication. Children that do experience discomfort usually do fine with Tylenol or Ibuprofen (follow directions on the bottle).
EXTRACTION (TOOTH WIGGLE)	<ul style="list-style-type: none"> ◆ You may resume brushing tonight, brushing gently in the site of the extraction. ◆Starting tomorrow you may use a warm salt water rinses to keep the extraction site clean. Simply place 1 to 2 teaspoons of salt in a warm cup of water - stir and then have your child swish by mouth 3 to 6 times daily, or at least after each meal or snack. ◆ If your child is too young to swish, simply dip a wash cloth in a the mixture and swab the extraction site well after each feeding. This area should heal up in approximately 2 to 3 weeks. ◆Soft foods are usually tolerated the best for the first few days following removal of the tooth/teeth. ◆Do not allow your child to suck from a straw or drink carbonated beverages for 48 hours.
SPACE MAINTAINER	<ul style="list-style-type: none"> ◆If your child has received a space maintainer do not allow him/her to eat any sticky, gooey or caramel type candies. These types of items frequently dislodge the spacer and require that the appliance be recemented. To avoid such problems please prohibit your child from eating such items. Please note that some children may find the appliance uncomfortable at first but will get use to it with time.
PAIN	<ul style="list-style-type: none"> ◆If your child appears to be in any pain or discomfort give Tylenol, following the instructions on the back of the bottle. If the doctor feels that your child will need a medication stronger, then one will be prescribed. Please note that most children, even with extensive dental work, do very well post-operatively with regular Tylenol Elixir or Ibuprofen. Do not hesitate to call the doctor if you have any questions regarding medications.
WHAT TO EAT	<ul style="list-style-type: none"> ◆Your child should have his/her diet limited to soft foods for 2 hours following the dental procedure to prevent biting the lip or tongue while numb. The following foods are acceptable: milk shake, ice cream, pudding, jello, soup, mashed potatoes, noodles, yogurt or any other foods of this consistency.

